

A DATE OF DATA COLLECTION:

B TIME OF DATA COLLECTION:

C WARD WHERE DATA IS COLLECTED:

D RESEARCH ASSISTANT CODE:

PATIENT CODE (LOCAL):

ROOM NUMBER:

SCREENING - INCLUSION CRITERIA

	yes	no	
S1 Time of data collection is POD1 AND patient is 6 hrs (minimum) in the ward End surgery: Date: 2 0 2 Y M M D D Time: H H M M POD1? Back in ward: Date: 2 0 2 Y M M D D Time: H H M M 6HRS?	☐	☐	If yes to 1 and 2 and 3 • Give the Outcomes questionnaire to the patient • Complete the Process questionnaire
S2 Patient is consenting age or over	☐	☐	If no to 1 or 2 or 3: • Do not fill in the rest of the Process questionnaire • Do not give the Outcomes questionnaire to the patient • Input the screening data (up to the point you have reached) into the web mask
S3 Patient has given his assent (or consent) to participate If no to S3, mark the reason(s): ☐ a. Patient is not on the ward ☐ b. Patient does not wish to participate ¹ ☐ b1. too ill ☐ b2. too much pain ☐ b3. other ☐ c. Patient is asleep ☐ d. Patient has visitors ☐ e. It is not possible to communicate with the patient (e.g., patient is deaf, does not read/write in any of the languages in which the Outcomes questionnaire is available) ☐ f. Patient is cognitively impaired (e.g., Downs syndrome, dementia, Alzheimer's disease, Cerebral Palsy) ☐ g. Other, specify:	☐	☐	Special case: If yes to 1 and 2 and 3f and you have permission from the Ethics Committee in your hospital: • Complete the Process questionnaire • Do not give the Outcomes questionnaire to the patient

¹ Remember: You may interview patients who need help, e.g., are too ill or in too much pain or illiterate

DEMOGRAPHIC INFORMATION

D1 Gender ☐ Male ☐ Female	D2 Year of birth Y Y																																			
D3 Weight kg	D4 Height cm																																			
D5 Nationality (check records)	D6 Country of birth (check records)																																			
D7 Language of Outcome questionnaire (select one) <table border="0"> <tr> <td>☐ Albanian</td> <td>☐ Arabic</td> <td>☐ Afrikaans</td> <td>☐ Bahasa Malaysia</td> <td>☐ Braz. Portuguese</td> <td>☐ Bulgarian</td> <td>☐ Chinese (Simp.)</td> </tr> <tr> <td>☐ Chinese (Trad.)</td> <td>☐ Czech</td> <td>☐ Dutch</td> <td>☐ English</td> <td>☐ Filipino</td> <td>☐ Finnish</td> <td>☐ French</td> </tr> <tr> <td>☐ German</td> <td>☐ Hebrew</td> <td>☐ Hindi</td> <td>☐ Hungarian</td> <td>☐ Icelandic</td> <td>☐ Indonesian</td> <td>☐ Italian</td> </tr> <tr> <td>☐ Korean</td> <td>☐ Latvian</td> <td>☐ Norwegian</td> <td>☐ Romanian</td> <td>☐ Russian</td> <td>☐ Serbian</td> <td>☐ Setswana</td> </tr> <tr> <td>☐ Spanish</td> <td>☐ Span. Mexico</td> <td>☐ Swedish</td> <td>☐ Turkish</td> <td>☐ Ukrainian</td> <td>☐ Xhosa</td> <td>☐ Zulu</td> </tr> </table>		☐ Albanian	☐ Arabic	☐ Afrikaans	☐ Bahasa Malaysia	☐ Braz. Portuguese	☐ Bulgarian	☐ Chinese (Simp.)	☐ Chinese (Trad.)	☐ Czech	☐ Dutch	☐ English	☐ Filipino	☐ Finnish	☐ French	☐ German	☐ Hebrew	☐ Hindi	☐ Hungarian	☐ Icelandic	☐ Indonesian	☐ Italian	☐ Korean	☐ Latvian	☐ Norwegian	☐ Romanian	☐ Russian	☐ Serbian	☐ Setswana	☐ Spanish	☐ Span. Mexico	☐ Swedish	☐ Turkish	☐ Ukrainian	☐ Xhosa	☐ Zulu
☐ Albanian	☐ Arabic	☐ Afrikaans	☐ Bahasa Malaysia	☐ Braz. Portuguese	☐ Bulgarian	☐ Chinese (Simp.)																														
☐ Chinese (Trad.)	☐ Czech	☐ Dutch	☐ English	☐ Filipino	☐ Finnish	☐ French																														
☐ German	☐ Hebrew	☐ Hindi	☐ Hungarian	☐ Icelandic	☐ Indonesian	☐ Italian																														
☐ Korean	☐ Latvian	☐ Norwegian	☐ Romanian	☐ Russian	☐ Serbian	☐ Setswana																														
☐ Spanish	☐ Span. Mexico	☐ Swedish	☐ Turkish	☐ Ukrainian	☐ Xhosa	☐ Zulu																														

BLANK FIELDS

Blank field 1:	Blank field 5:
Blank field 2:	Blank field 6:
Blank field 3:	Blank field 7:
Blank field 4:	Project phase:

MEDICAL HISTORY**H1 Comorbidities**
☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

Cancer	☐ Cancer
Renal	☐ Renal insufficiency or disease without dialysis ☐ Renal disease requiring dialysis
Diabetes	☐ Diabetes Type I ☐ Diabetes Type II ☐ Diabetes Type unknown
Psychiatric	☐ Affective disorders (depression, anxiety, phobia, PTSD, bipolar disorder) ☐ Schizophrenia ☐ Alcohol use disorder ☐ Current smoker ☐ Substance abuse of drugs (legal and illegal)
Cardiovascular	☐ Hypertension ☐ Coronary artery disease or myocardial infarction or cerebral vascular accident
Hematology	☐ Sickle cell disease
GI disease	☐ Liver Cirrhosis ☐ History or current upper or lower GI ulcer (peptic or duodenal ulcer disease) ☐ Irritable bowel disease (Crohn's disease, ulcerative colitis)
Pulmonary disease	☐ Asthma ☐ Sleep apnea ☐ Chronic Obstructive Pulmonary Disease (COPD)
Neurologic	☐ Fibromyalgia
Steroid use	☐ Regular administration of oral or parenteral corticosteroid medications
Musculoskeletal	☐ Osteoarthritis ☐ Rheumatoid arthritis
Multiple trauma	☐ At least 1 fracture(s) / laceration(s) / tissue damage in addition to the current reason for surgery
Other surgery	☐ Patient has already undergone another surgery during current hospitalization
	☐ Other , specify:

H2 Existing condition (check medical record)
☐ Pregnancy, Week:
☐ not relevant ☐ not possible to obtain the information

☐ Lactation

☐ not relevant ☐ not possible to obtain the information
H3 Did the patient receive any opioid(s) before the current admission?
☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	Immediate release (PO & other)	Controlled release; (PO & other)
Buprenorphine	☐ mg\day	☐ µg\hr transdermal
Codeine	☐ mg\day	☐ mg\day
Fentanyl	☐ µg\hr transmucosal / intranasal	☐ µg\hr transdermal
Hydrocodone	☐ mg\day	☐ mg\day
Hydromorphone	☐ mg\day	☐ mg\day
Morphine	☐ mg\day	☐ mg\day
Oxycodone	☐ mg\day	☐ mg\day
Oxycodone (with Naloxon)	☐ mg\day	☐ mg\day
Pethidine (Meperidine)	☐ mg\day	☐ mg\day
Tapentadol	☐ mg\day	☐ mg\day
Tilidin (w\wo Naloxon)	☐ mg\day	☐ mg\day
Tramadol	☐ mg\day	☐ mg\day
Other, specify: 	☐	☐
Other, specify: 	☐	☐

PRE - MEDICATION

M1 Sedatives (pre-medication)

☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	p.o.	i.v.	i.m.
Diazepam	☐ mg	☐ mg	☐ mg
Clorazepate dipotassium	☐ mg	☐ mg	☐ mg
Haloperidol	☐ mg	☐ mg	☐ mg
Lorazepam	☐ mg	☐ mg	☐ mg
Midazolam	☐ mg	☐ mg	☐ mg
Promethazine	☐ mg	☐ mg	☐ mg
Other, specify: 	☐ mg	☐ mg	☐ mg

M2 Non-opioids (pre-medication)

☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	p.o.	i.v.	i.m.	supp.
Celecoxib	☐ mg	☐ mg	☐ mg	☐ mg
Dexamethasone	☐ mg	☐ mg	☐ mg	☐ mg
Dexketoprofen	☐ mg	☐ mg	☐ mg	☐ mg
Diclofenac	☐ mg	☐ mg	☐ mg	☐ mg
Etoricoxib	☐ mg	☐ mg	☐ mg	☐ mg
Flurbiprofen	☐ mg	☐ mg	☐ mg	☐ mg
Gabapentin	☐ mg	☐ mg	☐ mg	☐ mg
Ibuprofen	☐ mg	☐ mg	☐ mg	☐ mg
Ketoprofen	☐ mg	☐ mg	☐ mg	☐ mg
Ketorolac	☐ mg	☐ mg	☐ mg	☐ mg
Metamizol	☐ mg	☐ mg	☐ mg	☐ mg
Naproxen	☐ mg	☐ mg	☐ mg	☐ mg
Nefopam	☐ mg	☐ mg	☐ mg	☐ mg
Paracetamol (Acetaminophen)	☐ mg	☐ mg	☐ mg	☐ mg
Parecoxib	☐ mg	☐ mg	☐ mg	☐ mg
Pregabalin	☐ mg	☐ mg	☐ mg	☐ mg
Other, specify: 	☐	☐	☐	☐

PRE - MEDICATION

M3 Opioids (pre-medication)

☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	Immediate release (PO & other)	Controlled release (PO & other)	i.v.	i.m.	supp.	s.c.
Buprenorphine	☐ mg	☐ µg/hr	☐ µg	☐ µg	☐ µg	☐ µg
Codeine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Dezocine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Fentanyl	☐ µg transmucosal	☐ µg/hr transdermal	☐ µg	☐ µg	☐ µg	☐ µg
Hydrocodone	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Hydromorphone	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Morphine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Nalbuphine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Oxycodone	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Oxycodone (with Naloxon)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Pethidine (Meperidine)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Piritramide	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Sufentanil	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg
Tapentadol	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Tilidin (w/wo Naloxon)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Tramadol	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg
Other, specify: 	☐	☐	☐	☐	☐	☐

SURGICAL PROCEDURE(S)

P1 Surgical procedure(s)

use ICD-9 codes link <http://icd9cm.chrisendres.com/index.php?action=proclist>

ICD-9 Procedure Code		Text (only for your notes, not necessary for mask)	
1		1	
2		2	
3		3	
4		4	

P2 Duration of surgery

Start surgery:

Date: 2 0 1 Y M M D DTime: H H M M

End surgery:

Date: 2 0 1 Y M M D DTime: H H M M

INTRA-OPERATIVE

M4 General anaesthesia (intra-op)
☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

☐ Inhalational	☐ IV
---------------------------------------	-----------------------------

M5 Regional anaesthesia (RA) (intra-op)
☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

☐ Epidural	☐ Spinal	☐ Brachial plexus	☐ Femoral
☐ Sciatic	☐ Paravertebral	☐ Transv. Abdom. Plane (TAP)	☐ Other:

In M8: Mark the RA medication(s) given in the RA column

M6 Non-opioids (intra-op)
☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	i.v.	i.m.	supp.
Clonidine	☐ mg	☐ mg	☐ mg
Dexamethasone	☐ mg	☐ mg	☐ mg
Dexketoprofen	☐ mg	☐ mg	☐ mg
Dexmedetomidine	☐ mg	☐ mg	☐ mg
Diclofenac	☐ mg	☐ mg	☐ mg
Flurbiprofen	☐ mg	☐ mg	☐ mg
Ibuprofen	☐ mg	☐ mg	☐ mg
Ketamine	☐ mg	☐ mg	☐ mg
Ketoprofen	☐ mg	☐ mg	☐ mg
Ketorolac	☐ mg	☐ mg	☐ mg
Metamizol	☐ mg	☐ mg	☐ mg
Naproxen	☐ mg	☐ mg	☐ mg
Nefopam	☐ mg	☐ mg	☐ mg
Paracetamol (Acetaminophen)	☐ mg	☐ mg	☐ mg
Parecoxib	☐ mg	☐ mg	☐ mg
Other, specify: 	☐	☐	☐
Other, specify: 	☐	☐	☐
	i.v.	i.m.	supp.

INTRA-OP

M7 Wound infiltration (intra-op)

☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible; analgesic is not recorded):

☐ Single shot by surgeon ☐ Indwelling catheter ☐ Other, specify: ☐ Other, specify:

M8 Opioids & local anaesthetics (intra-op)

☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	RA (see M5)	i.v.	i.m.	s.c.
Alfentanil	☐ mg	☐ mg	☐ mg	☐ mg
Buprenorphine	☐ µg	☐ µg	☐ µg	☐ µg
Codeine	☐ mg	☐ mg	☐ mg	☐ mg
Dezocine	☐ mg	☐ mg	☐ mg	☐ mg
Fentanyl	☐ µg	☐ µg	☐ µg	☐ µg
Hydrocodone	☐ mg	☐ mg	☐ mg	☐ mg
Hydromorphone	☐ mg	☐ mg	☐ mg	☐ mg
Morphine	☐ mg	☐ mg	☐ mg	☐ mg
Nalbuphin	☐ mg	☐ mg	☐ mg	☐ mg
Oxycodone	☐ mg	☐ mg	☐ mg	☐ mg
Pethidine (Meperidine)	☐ mg	☐ mg	☐ mg	☐ mg
Piritramid	☐ mg	☐ mg	☐ mg	☐ mg
Remifentanyl	☐ mg	☐ mg	☐ mg	☐ mg
Sufentanyl	☐ µg	☐ µg	☐ µg	☐ µg
Tramadol	☐ mg	☐ mg	☐ mg	☐ mg
Bupivacaine	☐	☐	☐	☐
Levobupivacaine	☐	☐	☐	☐
Lidocaine	☐	☐	☐	☐
Prilocaine	☐	☐	☐	☐
Ropivacaine	☐	☐	☐	☐
Other, specify: 	☐	☐	☐	☐
Other, specify: 	☐	☐	☐	☐
	RA	i.v.	i.m.	s.c.

RECOVERY ROOM**M9 Non-opioids (recovery room)**
☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	p.o.	i.v.	i.m.	supp.
Celecoxib	☐ mg	☐ mg	☐ mg	☐ mg
Clonidine	☐ mg	☐ mg	☐ mg	☐ mg
Dexamethasone	☐ mg	☐ mg	☐ mg	☐ mg
Dexketoprofen	☐ mg	☐ mg	☐ mg	☐ mg
Diclofenac	☐ mg	☐ mg	☐ mg	☐ mg
Etoricoxib	☐ mg	☐ mg	☐ mg	☐ mg
Flurbiprofen	☐ mg	☐ mg	☐ mg	
Gabapentin	☐ mg	☐ mg	☐ mg	☐ mg
Ibuprofen	☐ mg	☐ mg	☐ mg	☐ mg
Ketamine	☐ mg	☐ mg	☐ mg	☐ mg
Ketoprofen	☐ mg	☐ mg	☐ mg	☐ mg
Ketorolac	☐ mg	☐ mg	☐ mg	☐ mg
Metamizol	☐ mg	☐ mg	☐ mg	☐ mg
Naproxen	☐ mg	☐ mg	☐ mg	☐ mg
Nefopam	☐ mg	☐ mg	☐ mg	☐ mg
Paracetamol (Acetaminophen)	☐ mg	☐ mg	☐ mg	☐ mg
Parecoxib	☐ mg	☐ mg	☐ mg	☐ mg
Pregabalin	☐ mg	☐ mg	☐ mg	☐ mg
Other, specify: 	☐	☐	☐	☐
Other, specify: 	☐	☐	☐	☐

M10 Regional analgesia (recovery room)
☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

☐ Epidural	☐ Spinal	☐ Brachial plexus	☐ Femoral
☐ Sciatic	☐ Paravertebral	☐ Transv. Abdom. Plane (TAP)	☐ Other:

In M11: (1) Mark the RA medication(s) given in the RA column

(2) If the medication was given as PCA, tick appropriate box in the PCA column

RECOVERY ROOM

M11 Opioids & local anaesthetics (recovery room)

☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	Immediate release (PO & other)	Controlled release (PO & other)	RA (see M10)	i.v.	i.m.	supp.	s.c.	PCA (see M10)
Buprenorphine	☐ mg	☐ µg/hr	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg	☐
Codeine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Dezocine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Fentanyl	☐ µg transmucosal	☐ µg/hr transdermal	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg	☐
Hydrocodone	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Hydromorphone	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Morphine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Nalbuphin	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Oxycodone	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Oxycodone (with Naloxone)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Pethidine (Meperidine)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Piritramid	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Sufentanil	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg	☐
Tapentadol	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Tilidin (w/o Naloxon)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Tramadol	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Bupivacaine	☐	☐	☐	☐	☐	☐	☐	☐
Levobupivacaine	☐	☐	☐	☐	☐	☐	☐	☐
Lidocaine	☐	☐	☐	☐	☐	☐	☐	☐
Prilocaine	☐	☐	☐	☐	☐	☐	☐	☐
Ropivacaine	☐	☐	☐	☐	☐	☐	☐	☐
Other, specify: 	☐	☐	☐	☐	☐	☐	☐	☐
Other, specify: 	☐	☐	☐	☐	☐	☐	☐	☐
Naloxone (only as an antagonist for respiratory depression)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
	Immediate release (PO & other)	Controlled release (PO & other)	RA	i.v.	i.m.	supp.	s.c.	PCA

WARD

M12 Non-opioids (ward)

☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	p.o.	i.v.	i.m.	supp.
Celecoxib	☐ mg	☐ mg	☐ mg	☐ mg
Clonidine	☐ mg	☐ mg	☐ mg	☐ mg
Dexamethasone	☐ mg	☐ mg	☐ mg	☐ mg
Dexketoprofen	☐ mg	☐ mg	☐ mg	☐ mg
Diclofenac	☐ mg	☐ mg	☐ mg	☐ mg
Etoricoxib	☐ mg	☐ mg	☐ mg	☐ mg
Flurbiprofen	☐ mg	☐ mg	☐ mg	☐ mg
Gabapentin	☐ mg	☐ mg	☐ mg	☐ mg
Ibuprofen	☐ mg	☐ mg	☐ mg	☐ mg
Ketamine	☐ mg	☐ mg	☐ mg	☐ mg
Ketoprofen	☐ mg	☐ mg	☐ mg	☐ mg
Ketorolac	☐ mg	☐ mg	☐ mg	☐ mg
Metamizol	☐ mg	☐ mg	☐ mg	☐ mg
Naproxen	☐ mg	☐ mg	☐ mg	☐ mg
Nefopam	☐ mg	☐ mg	☐ mg	☐ mg
Paracetamol (Acetaminophen)	☐ mg	☐ mg	☐ mg	☐ mg
Parecoxib	☐ mg	☐ mg	☐ mg	☐ mg
Pregabalin	☐ mg	☐ mg	☐ mg	☐ mg
Other, specify: 	☐	☐	☐	☐

M13 Regional analgesia (ward)

☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

☐ Epidural	☐ Spinal	☐ Brachial plexus	☐ Femoral
☐ Sciatic	☐ Paravertebral	☐ Transv. Abdom. Plane (TAP)	☐ Other:

In M14: (1) Mark the RA medication(s) given in the RA column

(2) If the medication was given as PCA, tick appropriate box in the PCA column

WARD

M14 Opioids & local anaesthetics (ward)

☐ yes ☐ no ☐ not possible to obtain the information

If yes, which (multiple answers possible):

	Immediate release (PO & other)	Controlled release (PO & other)	RA (see M13)	i.v.	i.m.	supp.	s.c.	PCA (see M13)
Buprenorphine	☐ mg	☐ µg/hr	☐ µg	☐ µg	☐ µg	☐ µg	☐	☐
Codeine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Dezocine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Fentanyl	☐ µg transmucosal	☐ µg/hr transdermal	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg ☐ µg	☐
Hydrocodone	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Hydromorphone	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Morphine	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Nalbuphin	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Oxycodone	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Oxycodone (with Naloxone)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Pethidine (Meperidine)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Piritramid	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Sufentanil	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg	☐ µg	☐
Sufentanil sublin- gual (Zalviso)		☐ µg						☐
Tapentadol	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Tilidin (w/wo Naloxon)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Tramadol	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐
Bupivacaine	☐	☐	☐	☐	☐	☐	☐	☐
Levobupivacaine	☐	☐	☐	☐	☐	☐	☐	☐
Lidocaine	☐	☐	☐	☐	☐	☐	☐	☐
Prilocaine	☐	☐	☐	☐	☐	☐	☐	☐
Ropivacaine	☐	☐	☐	☐	☐	☐	☐	☐
Other, specify: 	☐	☐	☐	☐	☐	☐	☐	☐
Naloxone (only as an antagonist for respiratory depression)	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐ mg	☐

M15 Measurement of pain:

a) Since the patient returned from surgery, how many times was pain assessed and this was recorded?

☐ 0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ >9 times ☐ not possible to obtain the information

b) After treatment with an analgesic, was the pain re-assessed within 60 minutes?

☐ no analgesic was given
☐ 0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ >9 times ☐ not possible to obtain the information